



Thorius narismagnus (Amphibia: Plethodontidae): rediscovery at the type locality and detection of a new population

¹José L. Aguilar-López, ^{2,*}Paulina García-Bañuelos, ³Eduardo Pineda, and ⁴Sean M. Rovito

^{1,2,3}Red de Biología y Conservación de Vertebrados, Instituto de Ecología, A.C., Carretera antigua a Coatepec 351, El Haya, Xalapa, Veracruz, MEXICO ⁴Unidad de Genómica Avanzada (Langebio), Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, km 9.6 Libramiento Norte Carretera Irapuato-León, Irapuato, Guanajuato CP 36824, MEXICO

Abstract.—Of the 42 Critically Endangered species of plethodontid salamanders that occur in Mexico, thirteen have not been reported in more than ten years. Given the lack of reports since 1976, the minute plethodontid salamander *Thorius narismagnus* is widely considered as missing. However, this report describes the rediscovery of this minute salamander at the type locality (Volcán San Martín), as well as a new locality on Volcán Santa Marta, 28 km southeast of its previously known distribution, both in the Los Tuxtlas region of Veracruz, Mexico. The localities where *T. narismagnus* has been found are mature forests in a community reserve on Volcán San Martín and a private reserve on Volcán Santa Marta. The presence of maxillary teeth, generally absent in *Thorius*, are reported here in some *T. narismagnus* females. Two efforts which may contribute to the conservation of *Thorius narismagnus* are the preservation of the cloud forests where this species persists, as well as the determination of the presence and possible effect of chytrid fungus in these populations.

Keywords. Conservation, ecological reserve, Los Tuxtlas, missing species, minute salamander, molecular analysis, Mexico

Citation: Aguilar-López JL, García-Bañuelos P, Pineda E, Rovito SM. 2019. *Thorius narismagnus* (Amphibia: Plethodontidae): rediscovery at the type locality and detection of a new population. *Amphibian & Reptile Conservation* 13(2) [General Section]: 126–132 (e193).

Copyright: © 2019 Aguilar-López et al. This is an open access article distributed under the terms of the Creative Commons Attribution License [Attribution 4.0 International (CC BY 4.0): <https://creativecommons.org/licenses/by/4.0/>], which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. The official and authorized publication credit sources, which will be duly enforced, are as follows: official journal title *Amphibian & Reptile Conservation*; official journal website: amphibian-reptile-conservation.org.

Received: 1 October 2018; **Accepted:** 4 August 2019; **Published:** 8 November 2019.

Introduction

In Mexico, 132 species of plethodontid salamanders have been recorded (AmphibiaWeb 2019), 42 of which are Critically Endangered (CR) according to the International Union for the Conservation of Nature (IUCN 2019). This conservation status may be assigned due to various combinations of biological characteristics (e.g., restricted distribution or specific environmental requirements) and risk factors (e.g., habitat modification or climate change; Stuart et al. 2008). In recent decades, population declines have been observed or estimated for several species (Frias-Alvarez et al. 2010) and in some cases, despite intensive search efforts, finding them has not been possible.

According to the IUCN (2019), 13 Critically Endangered plethodontid species in Mexico have not been reported in more than 10 years. Since 2010, six of these species have been rediscovered: *Isthmura naucampatepetl* (Naturalista 2019), *Chiropterotriton magnipes*, *C. mosaueri*, *Pseudoeurycea ahuitzotl*, *P. tlahcuiloh*, and

Thorius munificus (AmphibiaWeb 2019). However, so far there have been no subsequent records of the remaining seven species, including *Thorius narismagnus*. The unknown status of these species highlights the need to carry out sampling efforts in their historical localities and to explore those areas with favorable environmental conditions, in order to locate new populations (Sandoval-Comte et al. 2012).

Thorius narismagnus (Shannon and Werler 1955) is a minute salamander with a known distribution that comprises only four localities, not more than 8 km apart, on Volcán San Martín in the Los Tuxtlas region, Veracruz, Mexico (Fig. 1), in an elevational range between 890 and 1,200 m asl (Hanken and Wake 1998). *Thorius narismagnus* is considered as CR because the extent of its occurrence is < 17 km² with reductions in the extent and quality of its habitat, and due to a continuing decline in the number of mature individuals (IUCN 2016). The historical records of *T. narismagnus* include 55 specimens collected between 1953 and 1976, and since

Correspondence. ¹jlal.herp@gmail.com; ^{2,*}paulinabanuelos@hotmail.es; ³eduardo.pineda@inecol.mx;
⁴sean.rovito@cinvestav.mx




Fig. 1. Locations of historical collection localities and the new localities of *Thorius narismagnus* in Los Tuxtlas region, Mexico.

then this species has not been reported (IUCN 2016).

Rediscovery

As part of a study on the diversity and conservation of amphibians in Veracruz, fieldwork was carried out in a cloud forest in the community reserve Ejido Ruiz Cortines, San Andrés Tuxtla, Veracruz (18°32'53"N, 95°09'16"W; 1,136 m asl) on Volcán San Martín (Fig. 1; Fig. 2A) and in the private Ecological Reserve "La Otra Opción" Catemaco, Veracruz (18°22'32"N, 94°55'28"W; 1,075 m asl) on Volcán Santa Marta (Fig. 1; Fig. 2B). At both sites searches for amphibians (08:00–12:00 and 20:00–00:00 h) were conducted in all terrestrial microhabitats commonly used by these organisms (Crump and Scott 1994).

In the Ejido Ruiz Cortines community reserve, with a cumulative search effort of 72 person-hours in September 2012, three individuals of the genus *Thorius* were detected, two of which were collected (from which a tissue sample was taken and the individuals were subsequently preserved) and deposited in the Colección de Anfibios y Reptiles del Instituto de Ecología A.C. (CARIE 0857, 1137; Fig. 2C). A sampling effort of 24 person-hours was carried out in July 2015 at this locality, but there were no sightings of any minute salamanders. The two collected specimens measured 17.6 and 11.4

mm in SVL, the ratios between the length and width of the nostrils were 1.14 and 1.09, with four and five free intercostal grooves separating the adpressed fore and hind limbs, respectively.

In the private ecological reserve La Otra Opción, with cumulative search efforts of 96 person-hours in July 2015 and 66 person-hours in July 2017, three individuals of *Thorius* were detected in well-preserved forests in each year. The three individuals detected in 2015 were captured (each one was measured and a sample of tail tissue was taken) and subsequently released; these specimens had SVL measurements of 15.4, 15.9, and 19.4 mm. The three individuals found in 2017 were collected (CARIE 1251, 1258, 1259; Fig. 2D) and a sample of tissue was taken from each; they measured 19.9, 22.5, and 21.2 mm SVL, respectively. The proportions between the length and width of the nostrils were 1, 1, and 1.15, respectively, with 5.5 free intercostal grooves separating adpressed limbs. Two females from La Otra Opción (CARIE 1251 and 1259) had eight and five maxillary teeth, respectively, while the adult male (CARIE 1258) lacked maxillary teeth.

Identification of Specimens

The coloration in life of the specimens collected in the Volcán San Martín and Volcán Santa Marta localities



Fig. 2. Habitat in the Volcán San Martín locality (A) and a specimen (C) collected from it in life (CARIE 0857). Habitat in the Volcán Santa Marta locality (B) and a specimen (D) collected from it in life (CARIE 1251).

was light brown with a dark brown spike pattern in the dorsum, dark brown color on the sides and a dark venter with small white spots (Fig. 2C, 2D). With the exception of the presence of maxillary teeth in two females, the morphological characters and the coloration coincide with the diagnosis proposed by Shannon and Werler (1955) and Rovito et al. (2013) for *Thorius narismagnus*. Although maxillary teeth are absent in most species of *Thorius*, they are present in several species, including *T. smithi*, which is relatively closely related to *T. narismagnus*. Furthermore, at least two species (*T. grandis* and *T. omiltemi*) have maxillary teeth present only in females, and maxillary teeth are more common in females of *T. minydemus* than in males, which rarely have them (Hanken and Wake 1998; Hanken et al. 1999). Maxillary teeth were absent in a total of 18 specimens from the type locality of *T. narismagnus* on Volcán San Martín as reported by Shannon and Werler (1955) and Hanken and Wake (1998), suggesting that they do not occur in either sex at that locality.

In order to confirm that both populations belong to *Thorius narismagnus*, DNA was extracted from liver tissue of one specimen from Volcán San Martín (the type locality of *T. narismagnus*) and two specimens from Volcán Santa Marta using a salt extraction protocol. A

784 bp fragment of the cytochrome b gene was amplified using primers MVZ15 and MVZ16 (Moritz et al. 1992). PCR consisted of an initial denaturation step of 94 °C for 2 min, followed by 35 cycles of denaturation at 94 °C for 30 sec, annealing at 48 °C for 1 min, and extension at 72 °C for 1 min, with a final extension at 72 °C for 7 min. PCR products were purified using ExoSAP IT (USB Corporation, Cleveland, Ohio, USA) and sequenced using the BigDye v3.1 terminator cycle sequencing kit (Applied Biosystems, Foster City, California, USA) on an ABI 3730 capillary sequencer. Sequences were edited using Geneious v8.1.8 (BioMatters, Auckland, New Zealand), and sequences used in analysis were 750 bp long after removing low-quality bases. Sequences for other species of *Thorius* were obtained from GenBank and sequences were aligned using Muscle v.3.8 (Edgar 2004). The pairwise GTR distance between sequences from the two populations was calculated using PAUP v4.165 (Swofford 2003). Sequences are deposited in GenBank (Table 1).

The average pairwise divergence for cytochrome b (cytb) between the two populations was 1.4% (Table 1). This level of divergence is comparable to, or lower than, that seen between conspecific populations of various species of *Thorius*. Several species of *Thorius*

Table 1. Percentage of average pairwise divergence for cytochrome b between the populations of *Thorius* species in GenBank.

Species	Voucher	GenBank accession number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32			
1 <i>T. arboreus</i>	IBH222720	KC884119	-																																		
2 <i>T. aureus</i>	IBH22356	KC884065	11.2	-																																	
3 <i>T. boreas</i>	IBH22339	KC884066	11.4	8.7	-																																
4 <i>T. dubitus</i>	MCZA137386	KC884068	16.6	16.0	18.7	-																															
5 <i>T. grandis</i>	MZFC27550	KC884115	10.6	11.4	12.3	16.7	-																														
6 <i>T. insperatus</i>	IBH22901	KC884120	19.1	18.2	17.2	15.3	16.9	-																													
7 <i>T. longicaudus</i>	MCZA148744	KC884096	7.9	8.5	10.0	14.8	7.3	15.3	-																												
8 <i>T. lunaris</i>	IBH22341	KC884069	18.8	19.9	18.7	18.8	20.8	20.6	19.6	-																											
9 <i>T. magdougalli</i>	IBH22890	KC884072	8.1	8.8	11.6	14.5	9.5	17.5	8.9	20.8	-																										
10 <i>T. magnipes</i>	IBH22918	KC884122	18.5	18.4	17.8	13.7	17.6	15.2	18.8	18.2	16.9	-																									
11 <i>T. maxillabrochus</i>	MCZA148743	KC884105	18.1	20.7	21.2	15.3	19.2	18.2	19.6	21.8	16.0	15.7	-																								
12 <i>T. minutissimus</i>	IBH23011	DQ640021	13.1	12.4	13.5	17.5	15.1	18.4	12.5	18.1	12.8	17.1	22.8	-																							
13 <i>T. minydemus</i>	MVZ229269	KC884100	23.3	28.4	28.7	17.5	22.9	19.1	24.7	22.2	23.0	16.3	8.4	24.0	-																						
14 <i>T. munificus</i>	IBH29716	KC884073	19.1	19.2	19.3	19.8	23.1	21.8	18.6	5.6	20.6	19.8	19.2	20.6	21.6	-																					
15 <i>T. narismagnus</i>	CARIE1251	MK761019	19.4	18.7	20.0	16.3	19.3	19.1	18.3	20.9	18.1	17.7	19.9	20.7	22.8	22.2	-																				
16 <i>T. narismagnus</i>	CARIE1258	MK761018	19.4	18.7	20.0	16.3	19.3	19.1	18.3	20.9	18.1	17.7	19.9	20.7	22.8	22.2	0.0	-																			
17 <i>T. narismagnus</i>	CARIE0857	MK761017	19.8	19.4	21.2	17.2	19.5	19.4	18.2	20.9	18.8	17.6	19.8	21.7	22.2	22.1	1.4	1.4	-																		
18 <i>T. narisovalis</i>	IBH26500	KC884077	12.2	10.3	11.0	18.6	14.6	15.6	9.4	19.1	11.4	17.3	19.1	12.1	22.7	19.8	18.9	18.9	18.6	-																	
19 <i>T. onilemi</i>	MVZ269309	KC884079	11.2	11.2	12.6	17.5	7.3	17.9	7.3	22.9	10.2	18.0	18.3	16.5	21.9	22.1	18.6	18.6	18.5	14.1	-																
20 <i>T. papaloae</i>	IBH22355	KC884081	10.7	10.0	11.1	17.4	11.4	16.5	9.0	19.7	9.3	17.2	18.8	16.1	22.4	20.0	18.8	18.8	18.5	10.8	10.1	-															
21 <i>T. pennatulus</i>	IBH26499	KC884082	20.4	20.7	20.5	18.5	19.9	20.2	18.5	20.1	20.0	18.0	18.9	21.1	18.8	19.8	8.8	8.8	8.6	22.1	18.4	19.9	-														
22 <i>T. pinicola</i>	IBH13995	KC884094	11.9	8.5	10.9	15.8	8.3	16.8	6.0	19.4	11.5	16.9	18.9	13.6	21.9	18.9	18.5	18.4	11.3	6.3	10.9	18.5	-														
23 <i>T. pulmonaris</i>	MCZA148742	KC884101	8.8	10.6	12.5	18.8	6.6	14.8	6.5	15.5	11.4	18.9	18.9	14.4	20.4	15.0	20.1	20.1	20.1	10.7	7.2	7.9	20.3	4.6	-												
24 <i>T. schmidtii</i>	MVZ269312	KC884085	19.3	21.0	20.0	12.1	19.8	18.3	19.7	22.6	19.8	9.2	15.6	20.5	17.3	22.4	17.7	17.7	18.6	19.5	21.2	18.9	18.9	18.7	20.9	-											
25 <i>T. smithi</i>	IBH26615	KC884121	19.6	18.6	18.7	15.8	20.9	13.6	16.9	21.2	18.2	17.0	14.9	19.8	16.6	20.0	10.3	10.3	10.2	17.5	18.5	16.6	11.8	17.7	18.5	17.4	-										
26 <i>T. sp. 1</i>	GP0099	KC884086	21.1	21.6	22.7	15.6	18.7	15.2	19.4	25.1	17.8	17.7	6.0	21.2	4.5	24.1	19.3	19.3	19.1	18.6	18.2	18.0	17.6	17.9	18.2	18.0	15.4	-									
27 <i>T. sp. 2</i>	MCZA148757	KC884118	9.6	9.4	10.4	17.2	8.1	16.2	6.2	17.9	9.6	18.6	19.9	13.1	22.3	18.3	18.4	18.4	18.3	12.4	7.5	9.9	17.7	6.9	6.6	22.6	18.1	18.8	-								
28 <i>T. sp. 3</i>	MCZA148759	KC884116	9.3	7.9	12.7	14.0	8.7	16.8	5.7	22.9	8.0	16.7	22.8	11.5	31.3	24.6	18.4	18.4	18.4	10.7	7.8	8.7	17.7	6.2	9.0	17.3	19.9	15.9	7.8	-							
29 <i>T. sp. 7</i>	MCZA148754	KC884111	13.2	11.7	13.9	19.7	14.4	18.5	11.3	22.4	9.8	22.4	19.7	18.1	25.3	22.0	20.6	20.6	21.0	13.7	13.2	9.3	22.5	12.7	13.2	21.9	19.2	17.7	12.4	10.2	-						
30 <i>T. spilogaster</i>	IBH22975	KC884087	18.5	20.3	21.1	14.2	18.2	14.4	18.8	20.5	17.5	14.8	7.4	20.0	2.3	20.8	17.7	17.7	18.1	18.5	18.0	17.0	16.2	17.2	17.3	15.6	14.5	4.5	18.0	14.2	17.6	-					
31 <i>T. tlaxiacus</i>	MCZA148745	KC884098	9.8	10.6	12.0	16.0	7.3	16.0	5.0	19.5	9.0	16.2	18.6	13.7	23.2	19.0	17.7	17.7	17.7	12.5	6.5	10.3	17.3	5.9	6.8	18.8	19.0	19.0	6.6	5.1	13.3	17.3	-				
32 <i>T. troglodytes</i>	IBH22981	KC614433	18.7	19.7	19.6	12.9	17.4	17.8	16.4	20.4	16.2	13.9	16.7	19.0	17.0	20.4	15.4	15.4	15.8	18.7	16.4	18.3	16.4	16.0	19.0	14.8	14.3	16.8	17.4	17.2	20.5	14.2	16.6	-			

from Oaxaca, including *T. boreas*, *T. macdougalli*, and *T. narisovalis*, have substantially higher divergence than that seen between the two populations from Los Tuxtlas that were sequenced here. For example, two populations of *Thorius boreas* separated by only 17 km are 5% divergent for *cytb* (Rovito et al. 2013). The close genetic similarity between the populations from Volcán Santa Marta and from the type locality of Volcán San Martín strongly suggests that these two populations are conspecific.

Conservation Implications

The record from the Ejido Ruiz Cortines community reserve locality represents the rediscovery of *T. narismagnus*, 36 years after the last reported record (Hanken 1976: MVZ183028–183035) at the type locality on Volcán San Martín (Hanken and Wake 1998). Additionally, the record from Volcán Santa Marta extends the distribution range of this species 28 km southeast of the closest known locality (Fig. 1). Because the specimens reported here were found in primary vegetation, and Díaz-García et al. (2017) recorded *T. narismagnus* in mature forest but not in restoration areas and cattle pasture in “La Otra Opción” this species is probably not able to survive in disturbed forest or even in moderately disturbed forest. Of the other missing species that the authors have found recently, only one (*T. munificus*) was found in small, highly disturbed forest fragments and within San Juan del Monte state reserve (Juárez-Ramírez et al. 2016), while *Chiropterotriton magnipes* and *C. mosaueri* were found in a cave in a national park with only light to moderate habitat disturbance, and *Pseudoeurycea ahuitzotl* and *P. tlahcuiloh* were found in intact montane forest (AmphibiaWeb 2019).

To more fully understand the conservation status of *T. narismagnus*, an exhaustive sampling effort through time is needed to determine how the encounter rate of this species varies throughout the year and whether it is currently an uncommon species, and to obtain a more accurate estimate of the population size. Extensive fieldwork is also necessary in those areas with favorable environmental conditions on both volcanoes to determine the full distribution of this species. The extent of well-preserved forest on Volcán San Martín is ~100 km², while on Volcán Santa Marta it is ~185 km² (INEGI 2016).

In addition, determining the presence of the chytrid fungus *Batrachochytrium dendrobatidis* (*Bd*) and its possible effect on the survival of these *T. narismagnus* populations is critical, because the presence of *Bd* in Los Tuxtlas region has been confirmed (Mendoza-Almeralla et al. 2015) and this pathogen is suspected of being linked to the population declines of this species (IUCN 2016). The presence of *Bd* has been reported in geographically close species from Central Veracruz, such as *Bolitoglossa rufescens*, *Aquiloerycea cephalica*, and *P. firscheini* (Van Rooij et al. 2011), as well as *P. nigromaculata*

and *Thorius pennatulus*, for which the presence of the pathogen has been associated with declines in their populations (Cheng et al. 2011).

Finally, the finding of two populations of a Critically Endangered salamander species that has gone unrecorded for almost four decades highlights the importance of community and private reserves for harboring species in imminent danger of extinction. Although both reserves are relatively small (less than 200 ha) compared to the Los Tuxtlas Biosphere Reserve in which they are located, most of the land of these reserves is conserved forest surrounded by modified environments. In that sense, this work shows the complementarity between the governmental and non-governmental reserves for the protection of species at risk of extinction (see García-Bañuelos et al. 2019), particularly those species with restricted distributions and sensitivity to environmental disturbances.

Acknowledgements.—We are grateful to Ricardo Luría, Luis Carrillo, Arístides García, Juan Díaz, David González, and Rogelio Agapito for fieldwork support; and to Ismael Guzmán for initial molecular analysis. Wesley Dáttilo provided helpful suggestions that improved this manuscript. Scientific collection permits were issued by Secretaría del Medio Ambiente y Recursos Naturales (SGPA/DGVS/03665/06 and SGPA/DGVS/03444/15).


Literature Cited

- AmphibiaWeb. 2019. AmphibiaWeb: Information on Amphibian Biology and Conservation. Berkeley, California, USA. Available: <http://amphibiaweb.org> [Accessed: 20 May 2019].
- Cheng TL, Rovito SM, Wake DB, Vredenburg VT. 2011. Coincident mass extirpation of neotropical amphibians with the emergence of the infectious fungal pathogen *Batrachochytrium dendrobatidis*. *Proceedings of the National Academy of Sciences of the United States of America* 108(23): 9,502–9,507.
- Crump ML, Scott Jr. NJ. 1994. Visual encounters surveys. Pp. 84–92 In: *Measuring and Monitoring Biological Diversity: Standard Methods for Amphibians*. Editors, Heyer RW, Donnelly MA, McDiarmid RW, Hayek LC, Foster MS. Smithsonian Institution Press, Washington, DC, USA. 384 p.
- Díaz-García JM, Pineda E, López-Barrera F, Moreno CE. 2017. Amphibian species and functional diversity as indicators of restoration success in tropical montane forest. *Biodiversity and Conservation* 26(11): 2,569–2,589.
- Edgar RC. 2004. MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics* 5(113): 1–19.
- Frias-Alvarez P, Zúñiga-Vega JJ, Flores-Villela O. 2010. A general assessment of the conservation status and decline trends of Mexican amphibians. *Biodiversity*


- and Conservation* 19(13): 3,699–3,742.
- García-Bañuelos P, Rovito SM, Pineda E. 2019. Representation of threatened biodiversity in protected areas and identification of complementary areas for their conservation: plethodontid salamanders in Mexico. *Tropical Conservation Science* 12: 1–15.
- Hanken J, Wake DB. 1998. Biology of tiny animals: systematics of the minute salamanders (*Thorius*: Plethodontidae) from Veracruz and Puebla, Mexico, with description of five new species. *Copeia* 1998(2): 312–345.
- Hanken J, Wake DB, Freeman H. 1999. Three new species of minute salamanders (*Thorius*: Plethodontidae) from Guerrero, Mexico, including the report of a novel dental polymorphism in Urodeles. *Copeia* 1999: 917–931.
- INEGI (Instituto Nacional de Estadística y Geografía). 2016. *Conjunto de datos vectoriales de Uso de Suelo y Vegetación escala 1:250 000, Serie VI (capa unión) (ed. 1)*. INEGI, Aguascalientes, Mexico.
- IUCN (International Union for Conservation of Nature). 2019. The IUCN Red List of Threatened Species. Version 2018-2. Available: <http://www.iucnredlist.org>. [Accessed: 11 June 2019].
- IUCN SSC Amphibian Specialist Group. 2016. *Thorius narismagnus*. The IUCN Red List of Threatened Species 2016: e.T59420A53986703. Available: <http://www.iucnredlist.org>. [Accessed: 6 March 2019].
- Juárez-Ramírez MC, Aguilar-López JL, Pineda E. 2016. Protected natural areas and the conservation of amphibians in a highly transformed mountainous region in Mexico. *Herpetological Conservation and Biology* 11(1): 19–28.
- Mendoza-Almeralla C, Burrowes P, Parra-Olea G. 2015. La quitridiomycosis en los anfibios de México. *Revista Mexicana de Biodiversidad* 86(1): 238–248.
- Moritz C, Schneider J, Wake DB. 1992. Evolutionary relationships within the *Ensatina eschscholtzii* complex confirm the ring species interpretation. *Systematic Biology* 41(3): 273–291.
- Naturalista. 2019. Salamandra de Cofre de Perote (*Isthmura naucampatepetl*). CONABIO (Comisión Nacional para el Conocimiento y Uso de la Biodiversidad), Mexico City, Mexico. Available: <https://www.naturalista.mx/taxa/476531-Isthmura-naucFampatepetl>. [Accessed: 6 March 2019].
- Rovito SM, Parra-Olea P, Hanken J, Bonett RM, Wake DB. 2013. Adaptive radiation in miniature: the minute salamanders of the Mexican highlands (Amphibia: Plethodontidae: *Thorius*). *Biological Journal of the Linnean Society* 109(3): 622–643.
- Sandoval-Comte A, Pineda E, Aguilar-López JL. 2012. In search of critically endangered species: the current situation of two tiny salamander species in the Neotropical mountains of Mexico. *PLoS ONE* 7(4): e34023.
- Shannon FA, Werler JE. 1955. Notes on amphibians of the Los Tuxtlas Range of Veracruz, Mexico. *Transactions of the Kansas Academy of Science* 58(3): 360–386.
- Stuart SN, Hoffmann M, Chanson JS, Cox NA, Berridge RJ, Ramani P, Young BE. 2008. *Threatened Amphibians of the World*. Lynx Ediciones, Barcelona, Spain; IUCN, Gland, Switzerland; and Conservation International, Arlington, Virginia, USA. 758 p.
- Swofford DL. 2003. *PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods)*. Version 4. Sinauer Associates, Sunderland, Massachusetts, USA.
- Van Rooij P, Martel A, Nerz J, Voitel S, Van Immerseel F, Haesebrouck F, Pasmans F. 2011. Detection of *Batrachochytrium dendrobatidis* in Mexican bolitoglossine salamanders using an optimal sampling protocol. *EcoHealth* 8(2): 237–243.




José L. Aguilar-López was born in Mexico, and obtained his Bachelor's degree at Benemérita Universidad Autónoma de Puebla (BUAP), and his M.Sc. degree at the Instituto de Ecología, A.C. (INECOL), both in Mexico. José recently obtained his Ph.D. degree at INECOL, and he is interested in the diversity, ecology, and conservation of amphibians and reptiles in tropical environments.



Paulina García-Bañuelos is a biologist, and recently obtained her Ph.D. degree at the Instituto de Ecología, A.C. (INECOL) in Xalapa, Veracruz, Mexico. Paulina obtained her M.Sc. degree at the Instituto de Neuroetología at the Universidad Veracruzana, the same university where she obtained her Bachelor's degree. She is interested in the current status and conservation of the plethodontid salamanders of Mexico.



Eduardo Pineda obtained his Bachelor's degree at the School of Sciences at the Universidad Autónoma del Estado de México (UAEM) and his doctoral degree at the Instituto de Ecología, A.C. (INECOL), in Xalapa, Mexico. Eduardo's research in the INECOL is focused on understanding the relationship between the transformation of tropical forests and biodiversity at different spatial scales, recognizing the importance of conserved areas and modified habitats for maintaining the diversity of amphibians, and assessing the current status through fieldwork of amphibian species that are in imminent danger of extinction. Currently, Eduardo has several undergraduate and graduate students addressing topics on the ecology and conservation of amphibians and reptiles in Mexico.



Sean Rovito is a professor at the National Laboratory of Genomics for Biodiversity (Langebio, Cinvestav) in Irapuato, Mexico. Sean's research focuses on diversification, genomics, and conservation of Neotropical salamanders, particularly the plethodontid salamanders of Mexico.